

List of Figures

Figure 1-1	SRS Organization	1-3
Figure 1-2	The Savannah River Site and Surrounding Area	1-7
Figure 3-1	D-Area Ash Project	3-4
Figure 3-2	Pathway for Processing and Dispositioning of Radioactive Liquid Waste at SRS.....	3-5
Figure 3-3	Salt Disposal Unit 6 (SDU-6)	3-7
Figure 4-1	Types and Typical Locations of Nonradiological Sampling.....	4-3
Figure 4-2	NPDES Industrial Wastewater Outfall Sampling Locations	4-5
Figure 4-3	NPDES Industrial Stormwater Outfall Sampling Locations.....	4-6
Figure 4-4	Nonradiological Surface Water Sampling Locations	4-8
Figure 4-5	Nonradiological Sediment Sampling Locations	4-10
Figure 4-6	Mercury in Sediment Locations.....	4-11
Figure 4-7	Mercury in Fish by Location and Species	4-12
Figure 5-1	Types and Typical Locations of Radiological Sampling.....	5-3
Figure 5-2	Ten-Year History of SRS Annual Tritium Releases to the Air	5-7
Figure 5-3	Percent of Tritium Released to the Air for 2015 and 2016	5-7
Figure 5-4	Air Sampling Locations Surrounding SRS up to 25 Miles.....	5-8
Figure 5-5	Radiological Liquid Effluent Sampling Locations	5-13
Figure 5-6	Ten-Year History of Direct Releases of Tritium to SRS Streams	5-14
Figure 5-7	Radiological Surface Water Sampling Locations	5-15
Figure 5-8	Ten-Year Trend of Tritium in Pen Branch and Fourmile Branch (pCi/L).....	5-18
Figure 5-9	Tritium Migration from SRS Seepage Basins and SWDF to SRS Streams	5-19
Figure 5-10	SRS Tritium Transport Summary	5-21
Figure 5-11	Offsite Drinking Water Sampling Locations.....	5-23
Figure 5-12	Tritium in Offsite Drinking Water and River Mile 118.8.....	5-24
Figure 5-13	Historical Trend of Average Cesium-137 Concentration in Deer Tissue (1965–2016).....	5-29
Figure 6-1	Exposure Pathways to Humans from Air and Liquid Effluents.....	6-4
Figure 6-2	2007–2011 Wind Rose Plot for H Area.....	6-5
Figure 6-3	Savannah River Annual Average Flow Rates at River Mile 118.8.....	6-6
Figure 6-4	Radionuclide Contributions to the 2016 SRS Total Liquid Pathway Dose of 0.15 mrem (0.0015 mSv).....	6-10
Figure 6-5	Radionuclide Contributions to the 2016 SRS Air Pathway Dose of 0.038 mrem (0.00038 mSv)	6-12

Figure 6-6	Ten-Year History of SRS Maximum Potential All-Pathway Doses.....	6-15
Figure 7-1	Groundwater at SRS.....	7-3
Figure 7-2	How Contamination Gets to Soil and Groundwater	7-4
Figure 7-3	Groundwater Plumes at SRS	7-6
Figure 7-4	Location of Site Boundary Wells at SRS—Between A/M Areas and Jackson, South Carolina	7-8
Figure 7-5	Location of Monitoring Wells Sampled for Tritium in Burke and Screven Counties, Georgia	7-11
Figure 7-6	Tritium Concentration in Wells Sampled in Burke and Screven Counties, Georgia	7-12
Figure 8-1	Interrelationship between QA/QC Activities	8-3